


BOOMERANG



A measurement by BOOMERANG of multiple peaks in the angular power spectrum of the cosmic microwave background

C.B. Netterfield¹, P.A.R. Ade², J.J. Bock³, J.R. Bond⁴, J. Borrill⁵, A. Boscaleri⁶, K. Coble⁷, C.R. Contaldi⁴, B.P. Crill⁸, P. de Bernardis⁹, P. Farese⁷, K. Ganga¹⁰, M. Giacometti⁹, E. Hivon¹⁰, V.V. Hristov⁸, A. Iacoangeli⁹, A.H. Jaffe¹¹, W.C. Jones⁸, A.E. Lange⁸, L. Martinis⁹, S. Masi⁹, P. Mason⁸, P.D. Mauskopf¹², A. Melchiorri¹³, T. Montroy⁷, E. Pascale⁶, F. Piacentini⁹, D. Pogosyan⁴, F. Pongetti⁹, S. Prunet⁴, G. Romeo¹⁴, J.E. Ruhl⁷, F. Scaramuzzi⁹

¹ *Depts. of Physics and Astronomy, University of Toronto, Canada*

² *Queen Mary and Westfield College, London, UK*

³ *Jet Propulsion Laboratory, Pasadena, CA, USA*

⁴ *Canadian Institute for Theoretical Astrophysics, University of Toronto, Canada*

⁵ *National Energy Research Scientific Computing Center, LBNL, Berkeley, CA, USA*

⁶ *IROE-CNR, Firenze, Italy*

⁷ *Dept. of Physics, Univ. of California, Santa Barbara, CA, USA*

⁸ *California Institute of Technology, Pasadena, CA, USA*

⁹ *Dipartimento di Fisica, Universita' La Sapienza, Roma, Italy*

¹⁰ *IPAC, California Institute of Technology, Pasadena, CA, USA*

¹¹ *Department of Astronomy, Space Sciences Lab and Center for Particle Astrophysics, University of CA, Berkeley, CA 94720 USA*


¹² *Dept. of Physics and Astronomy, Cardiff University, Cardiff CF24 3YB, Wales, UK*

¹³ *Nuclear and Astrophysics Laboratory, University of Oxford, Keble Road, Oxford, OX 3RH, UK*

¹⁴ *Istituto Nazionale di Geofisica, Roma, Italy*



ABSTRACT

This paper presents a measurement of the angular power spectrum of the Cosmic Microwave Background from $\ell = 75$ to $\ell = 1025$ ($\sim 10'$ to $5''$) from a combined analysis of four 150 GHz channels in the BOOMERANG experiment. The spectrum contains multiple peaks and minima, as predicted by standard adiabatic-inflationary models in which the primordial plasma undergoes acoustic oscillations. These results significantly constrain the values of Ω_{tot} , $\Omega_b h^2$, $\Omega_c h^2$ and n_s .





NASA balloon size

Boomerang Long-duration Cryostat and Telescope




Spiderweb Bolometers

- *small cosmic ray cross section*
- *reduced mass ->faster, stronger*



Launch Vehicle




BOOMERANG Launch




BOOMERANG Launch




BOOMERANG Launch




BOOMERANG Launch




ANTARCTIC Balloon Track




BALLOON at Float




ANTARCTIC Balloon Track




ANTARCIC Balloon Track



McMurdo Station, Antarctica


























CMB Results c. 2000–03–22

